

# Swan Canning Estuary Water Quality Monitoring Project

## Weekly Water Quality Report


### Canning Estuary and Lower Canning River

8 February 2022

Prepared by

Rivers and Estuaries Science  
Biodiversity and Conservation Science  
Department of Biodiversity, Conservation and Attractions

## Canning Estuary and Lower Canning River - Water Quality Profiles – 8 February 2022



Date: 8 February 2022

Weather & tide conditions: Conditions were slightly overcast with a north-westerly breeze of up to 4 knots. The predicted tides at Barrack St were 0.61 m at 6:55 am (low tide) and 1.0 m at 4:48 pm (high tide). Perth recorded 0.8 mm of rainfall in the week prior to sampling (Bureau of Meteorology).

Oxygenation: The Bacon St and Nicholson Rd oxygenation plants were triggered to provide oxygen in the 24 hours prior to sampling.

Canning Estuary (SCB2 to CASMID): The Canning Estuary was saline from SCB2 to CASMID. Surface waters were oxygenated from SCB2 to RIV and low in oxygen at CASMID. Bottom waters were low in oxygen at RIV and hypoxic at CASMID. Chlorophyll fluorescence was low throughout this zone. Water temperature ranged from 22.4 to 29.3 °C.

Lower Canning River (KEN to ELL): The Lower Canning River was fresh. Surface waters were oxygenated or well oxygenated and bottom waters were low in oxygen at ELL. Chlorophyll fluorescence was low throughout this zone. Water temperature ranged from 22.5 to 25 °C.

NB: Profile plots are visual interpolations of measured parameters only. Detailed data are available at [wir.water.wa.gov.au](http://wir.water.wa.gov.au).

**Oxygenation Plant Operational Status:**

- ▲ Operating for part or all of the 24 hours prior to sampling
- ▲ Operable but not triggered to operate in the 24 hours prior to sampling
- ▲ Inoperable for part or all of the 24 hours prior to sampling

**Definitions:**

Salinity – fresh <5, brackish 5-25, saline 25-35, hypersaline >35

Dissolved oxygen – well oxygenated >6 mg L<sup>-1</sup>, oxygenated >4-6 mg L<sup>-1</sup>, low oxygen >2-4 mg L<sup>-1</sup>, hypoxic 0.5-2 mg L<sup>-1</sup>, anoxic <0.5 mg L<sup>-1</sup>

Chlorophyll fluorescence (summer): low < 50 µg L<sup>-1</sup>, moderate 50-150 µg L<sup>-1</sup>, high 150-400 µg L<sup>-1</sup>, extreme > 400 µg L<sup>-1</sup>